

# ENVIRONMENTAL EXPOSURES AND HEALTH RISKS

Developed by  
The Environmental Health Sciences Research Center

Researched and Compiled by Lydia Johnston  
Questions? Visit our website [EHSRC.org](http://EHSRC.org)  
The EHSRC is funded by NIH P30 ES005605



# IOWA

---

College of Public Health

**The EHSRC would like to thank the development team (Lydia Johnson, Dr. Mary Charlton, Jackie Curnick, Dr. Brandi Janssen, and Dr. Hans Lehmler), and the expert reviewers who reviewed the content (Arsenic: Dr. Peter Thorne; Atrazine: Dr. Charles Lynch; Glyphosate: Dr. Charles Lynch; Nitrate: Dr. Mary Ward; and Radon: Dr. Jong Sung Kim).**



**View these resources online and check here for more fact sheets to be uploaded in the future.**

**<https://bit.ly/EHSRCResources>**

# Introduction



**IOWA**

College of Public Health

Iowa has the second-highest and fastest-growing rate of new cancer in the United States (1) as well as one of the greatest disparities in cancer mortality between Black and white populations (2). While many well-understood behavioral and genetic factors increase cancer risk, the role of exposures in the environment in increasing cancer risk is not as well documented. A chemical exposure can occur when a person's body comes into contact with a harmful substance in the air, water, or soil and has the potential to contribute to short-term and long-term illness.

A combination of natural and human activities contribute to environmental exposures to cancer-causing agents. Iowa has the largest percentage of homes that test at or above the EPA action level, as well as the highest average groundwater nitrate pollution from fertilizer use in the country. Thus, exposure to environmental carcinogens is an important consideration when examining the elevated cancer rates in the state (3, 4, 5). However, it should also be noted that higher than average red meat consumption, lower levels of activity, high alcohol consumption, UV exposure, some viruses, tobacco and nicotine exposure, lack of access to healthy fruits and vegetables, and more can further increase an individual's cancer risk, as highlighted by the Iowa Cancer Registry (1, 2). In line with priorities six and seven of Chapter 2 of the Iowa Cancer Plan, the goal of these fact sheets is to explain the current science about environmental contaminants and their relationship to cancer and other diseases to reduce the rates of potentially preventable diseases (6).

It is difficult to establish a causal relationship between one individual's exposure to an environmental carcinogen and cancer, so scientists use other ways to study these connections. In these fact sheets, risk, correlation, and association are all terms used to describe the relationship between an environmental contaminant and a health outcome (cancer, birth defects, etc.)

- **Risk:** explains the likelihood of an outcome in an exposed group compared to a lesser or unexposed group. A relative risk ratio of two indicates that the exposed group is two times more likely to experience the outcome than the comparison group.
- **Correlation:** measures the relationship between an exposure and a health outcome and how the two move together. A positive correlation indicates that when one variable increases, the other variable also increases. For example, when higher concentrations of radon are present in residential air, more people have lung cancer.
- **Association:** indicates there is a general relationship between an exposure and an outcome. A positive association means that an exposure is related to a higher probability of a health outcome.

In summary, relative risk ratio is the most specific indicator of a relationship, correlation shows a trend, and association is a more general relationship.

These fact sheets frequently reference Monographs produced by the International Agency for Research on Cancer (IARC). IARC evaluates carcinogenic risks by systematically reviewing and assessing studies published in peer-reviewed scientific literature. The evaluations consider evidence from human, animal, and mechanistic studies to ensure a rigorous and transparent review process. (8).

# References



**IOWA**

College of Public Health

1. Media Roundup: 2024 Cancer in Iowa Report. (2024). University of Iowa College of Public Health. Retrieved July 4, 2024, from <https://shri.public-health.uiowa.edu/wp-content/uploads/2024/02/cancer-in-iowa-2024.pdf>
2. Media Roundup: 2023 Cancer in Iowa Report. (2023). University of Iowa College of Public Health. Retrieved July 4, 2024, from <https://shri.public-health.uiowa.edu/wp-content/uploads/2023/02/cancer-in-iowa-2023.pdf>
3. Iowa Department of Health and Human Services (n.d.) Radon and You: What You Need to Know to Protect You and Your Family. <https://hhs.iowa.gov/media/7880/download?inline=>
4. Iowa Water Quality Information System (n.d.) Water Quality Parameters Explained. From July 4, 2024. <https://iwqis.iowawis.org/parameters.html#:~:text=It%20is%20an%20essential%20plant,tanks%2C%20and%20runoff%20from%20manure.>
5. Iowa Department of Natural Resources (n.d.). Iowa Source Water Protection Guidebook. From July 4, 2024. <https://www.iowadnr.gov/Portals/idnr/uploads/water/wse/SWPPGuidebook.pdf>
6. Iowa Cancer Consortium (n.d.). Iowa Cancer Plan – Prevention and Risk Reduction. Retrieved July 4, 2024, from <https://canceriowa.org/cancer-plan/introduction/#chapter-2>
7. United States Environmental Protection Agency (2024). Health Risk of Radon. <https://www.epa.gov/radon/health-risk-radon>
8. IARC. (2019). IARC Monographs on the identification of Carcinogenic Hazards to Humans: Preamble. <https://monographs.iarc.who.int/wp-content/uploads/2019/07/Preamble-2019.pdf>

# Arsenic & Health Fact sheet



# IOWA

College of Public Health

## Introduction

Inorganic arsenic is a toxic metalloid found naturally in Earth's crust. Arsenic can be detected in water, air, and soil globally and within the United States (1). While the metalloid can be found in many forms, inorganic arsenic is the environmental form that poses the greatest risk to public health (1). Arsenic is a colorless powder, with no smell or taste, making it imperceptible in water, food, air, and soil.

## Exposure Sources

Exposure to inorganic arsenic is associated with mining and smelting, proximity to hazardous waste sites, and its use as a pesticide, wood preservative, and alloy in semiconductors and batteries (1,2). While some exposure is attributable to the consumption of rice, the primary route of exposure to arsenic in the US is through drinking water where high concentrations of inorganic arsenic are present in the groundwater. Concentrations of inorganic arsenic may be high in groundwater used as a source of drinking water (3). Wells deeper than 100 feet have the highest risk of inorganic arsenic contamination.

## Health Risks Associated with Arsenic

### Cancer Health Risks

The U. S. Environmental Protection Agency (EPA) classifies inorganic arsenic compounds as carcinogenic to humans (4). Hundreds of epidemiological studies conducted worldwide have shown the exposure to arsenic causes cancers of the skin, bladder, and lung. There is also evidence of association with cancers of the kidney, nasal passages, liver, and prostate. Recent research in the U.S. has provided evidence of a significantly elevated risk at levels below the current drinking water standard of 10 micrograms per liter (7).

### Non-Cancer Health Risks

The EPA has conducted risk assessment on non-cancer effects of long-term exposure to inorganic arsenic including ischemic heart disease (IHD), Type II diabetes, adverse pregnancy and birth outcomes, and neurodevelopmental effects (7). Evidence demonstrates risk levels for IHD and Type 2 diabetes as high as 180 per 10,000 people with lifetime drinking water exposure to inorganic arsenic at the 10 micrograms per liter maximum contaminant level (MCL) (7).

## How is Arsenic regulated?

- The EPA established the MCL to be 10 micrograms per liter to protect against the cancer and non-cancer health effects listed above (5).
- Under the Safe Drinking Water Act, the EPA requires public water suppliers to test for inorganic arsenic on an ongoing basis(1,3).
- Private wells are unregulated for arsenic (1,3).
- The Food and Drug Administration (FDA) and Occupational Safety and Health Administration (OSHA) regulate dietary and occupational exposures, respectively (5).

**Published September 2024 -  
Questions? Visit our website  
EHSRC.org**

## What can you do?

- If your water is supplied from a private well, you should get your water tested annually for arsenic by contacting your local health department.
- If you are on a public water supply, read your annual Consumer Confidence Report to see what levels of inorganic arsenic are in your drinking water. Arsenic is not reported if the concentration is below 5 micrograms per liter (3).
- Certain methods can be used to enhance the removal of arsenic from drinking water. These include anion exchange, reverse osmosis, activated alumina, pH adjustment, enhanced lime softening, and pre-oxidation (3,6).
- Using a Brita filter, refrigerator filter, and boiling water are not reliable methods of arsenic removal (6).
- Rice should be rinsed thoroughly prior to cooking.

# References



**IOWA**

College of Public Health

1. United States Environmental Protection Agency (2001). Technical Fact Sheet: Final Rule for Arsenic in Drinking Water. <https://nepis.epa.gov/Exe/ZyPdf.cgi?Dockey=20001XXE.txt>
2. Centers for Disease Control and Prevention (2023). Routes of Exposure for Arsenic. Agency for Toxic Substances and Disease Registry. [https://www.atsdr.cdc.gov/csem/arsenic/what\\_routes.html](https://www.atsdr.cdc.gov/csem/arsenic/what_routes.html)
3. Moles, D. (2011) Arsenic in Iowa's Drinking Water. Iowa Department of Natural Resources. [https://www.iowadnr.gov/Portals/idnr/uploads/water/wells/Arsenic\\_information\\_sheet.pdf](https://www.iowadnr.gov/Portals/idnr/uploads/water/wells/Arsenic_information_sheet.pdf)
4. IARC Working Group on the Evaluation of Carcinogenic Risk to Humans (2012). IARC monographs on the evaluation of carcinogenic risk of chemicals to man. Arsenic, Metals, Fibres, and Dusts. IARC monographs on the evaluation of carcinogenic risk to humans, 100C.
5. Centers for Disease Control and Prevention (2023). What Are the Standards and Regulation for Arsenic Exposure? Agency for Toxic Substances and Disease Registry. <https://www.atsdr.cdc.gov/csem/arsenic/standards.html>
6. Barnaby, R., Liefeld, A., Jackson, B. P., Hampton, T. H., & Stanton, B. A. (2017). Effectiveness of table top water pitcher filters to remove arsenic from drinking water. Environmental research, 158, 610–615. <https://doi.org/10.1016/j.envres.2017.07.018>
7. Environmental Protection Agency. (2023). Peer Review of the IRIS Toxicological Review of Inorganic Arsenic. EPA. [https://sab.epa.gov/ords/sab/r/sab\\_apex/sab/advisoryactivitydetail?p18\\_id=2631&clear=18&session=10072872086921#doc](https://sab.epa.gov/ords/sab/r/sab_apex/sab/advisoryactivitydetail?p18_id=2631&clear=18&session=10072872086921#doc)

# Atrazine & Health Fact sheet



# IOWA

College of Public Health

## Introduction

Atrazine is a herbicide used to control broad-leaved weeds (1). Atrazine is used abundantly in Iowa, where it is applied to 65% of planted acres of corn (2). The herbicide is predominantly used in spring and summer (3). Atrazine is an odorless, white powder that dissolves in water; however, it breaks down slowly and persists in local water sources (4).

## Exposure Sources

People can be exposed to atrazine in many ways. While not detected in food sources, atrazine is found widely in lakes, rivers, estuaries and groundwater, leaving well water users at risk of exposure (1, 3). Most people are not exposed to atrazine regularly (1). However, well water users and those living on farms and in rural communities near areas where atrazine was applied to crops or manufactured may be exposed through contaminated dirt and water.(3,4). This exposure is particularly concerning for children who dig and play in outdoor spaces that may be contaminated. Further, occupational exposure from inhalation or absorption through the skin may occur (1).

## Health Risks Associated with Atrazine

### Cancer Health Risks

When last reviewed in 1999, the International Agency for Research on Cancer (IARC) indicated atrazine as “not classifiable” as to its human carcinogenicity, meaning there was inadequate evidence of human and/or animal carcinogenicity at the time. IARC will be releasing an updated evaluation in October of 2025. In an updated investigation of the Agricultural Health Study Cohort in 2024, atrazine exposure was associated with an increased risk of some cancers in farmers who were applying atrazine (5). Statistically significant associations were found for lung cancer, and for prostate cancer among those younger than sixty at diagnosis. Exposures to atrazine that occurred 25 years ago may lead to an elevated risk of pharyngeal and kidney cancer (5).

### Non-Cancer Health Risks

Exposure to atrazine is associated with a range of adverse health problems, primarily cardiovascular and reproductive system effects (6). The Environmental Protection Agency (EPA) established the Maximum Contaminant Level (MCL) to protect against adverse health effects to the cardiovascular system and reproductive problems associated with long-term exposure to atrazine at levels above the MCL. Some of the reproductive problems associated with this long-term exposure include an increased risk of pre-term delivery, low fetal weight, and heart, urinary, and limb defects in infants (1).

## How is Atrazine regulated?

- The EPA established the MCL for atrazine to be 0.003 mg/L in public drinking water to protect against cardiovascular or reproductive system effects (6).
- Atrazine is a restricted-use pesticide, meaning a person must be certified to purchase and apply this herbicide, due to its potential to contaminate the water and the environment (3).
- Public water suppliers are required to test for atrazine and present results to consumers and the state through consumer confidence reports (CCR). (Visit this link to find your local CCR: <https://ordspub.epa.gov/ords/safewater/f?p=136:102:::>)

## What can you do?

- Avoid areas for 12 hours following atrazine application. Do not allow children to play in soil and water sources following the application of atrazine (3, 4).
- If living near agricultural fields, ask your local health department about testing your private well for atrazine (7).
- Avoid tracking contaminated soil into homes, especially into rooms where children live and play. Vacuum floors and dust surfaces to avoid exposure to low levels of atrazine (3).
- Granular activated carbon filtration is an effective method to remove atrazine from contaminated water (8). These filtration systems can be applied over the whole house plumbing system, or as point of use filters just before the faucet.

Published March 2025  
Questions? Visit our website [EHSRC.org](http://EHSRC.org)

# References



**IOWA**

environmental health sciences  
— research center —

**College of Public Health**

1. DeMarini, D & Zahn, S. H. (1999) Atrazine. International Agency for Research on Cancer.
2. United States Department of Agriculture. (2021). Iowa Ag News—Chemical Use.  
[https://www.nass.usda.gov/Statistics\\_by\\_State/Iowa/Publications/Other\\_Surveys/2022/IA-Ag-Chem-Corn-2022.pdf](https://www.nass.usda.gov/Statistics_by_State/Iowa/Publications/Other_Surveys/2022/IA-Ag-Chem-Corn-2022.pdf)
3. Centers for Disease Control and Prevention. (n.d.) Atrazine: Public Health Statement. Agency for Toxic Substances and Disease Registry. <https://www.atsdr.cdc.gov/ToxProfiles/tp153-c1.pdf>
4. Centers for Disease Control and Prevention. (2011). ToxFAQs™ for Atrazine.  
<https://www.atsdr.cdc.gov/toxfaqs/tfacts153.pdf>
5. Remigio, R. V., Andreotti, G., Sandler, D. P., Erickson, P. A., Koutros, S., Albert, P. S., Hurwitz, L. M., Parks, C. G., Lubin, J. H., Hofmann, J. N., & Beane Freeman, L. E. (2024). An Updated Evaluation of Atrazine-Cancer Incidence Associations among Pesticide Applicators in the Agricultural Health Study Cohort. *Environmental health perspectives*, 132(2), 27010. <https://doi.org/10.1289/EHP13684>
6. Environmental Protection Agency. (2024). National Primary Drinking Water Regulations.  
<https://www.epa.gov/ground-water-and-drinking-water/national-primary-drinking-water-regulations>
7. Iowa Department of Natural Resources (2024). Private Well Program.  
<https://www.iowadnr.gov/environmental-protection/water-quality/private-well-program>
8. Oregon Health Authority. (2015). Atrazine and Drinking Water.  
<https://www.oregon.gov/oha/ph/healthyenvironments/drinkingwater/monitoring/documents/health/atrazine.pdf>



# Glyphosate & Health Fact sheet



# IOWA

College of Public Health

---

## Introduction

Glyphosate is a broad-spectrum herbicide, meaning it is toxic to a wide range of crop species. Glyphosate is used abundantly in agriculture to control the growth of broad-leaved weeds and grasses (1). Typically sold as Roundup or Ranger Pro, glyphosate is used in mixtures referred to as glyphosate-based herbicides (GBHs)(2,3). Not all Round Up contains glyphosate. To check whether a product contains glyphosate, check the product label. Glyphosate is colorless and odorless, making it imperceptible in the environment and on food, water, and dust(3). Due to the widespread use in agricultural and residential settings, exposure to glyphosate is very common. Glyphosate use increased substantially after 1996 with the innovation of glyphosate-resistant crops, which allowed the herbicide to be used throughout the growing season to manage weeds (3).

## Exposure Sources

Glyphosate residues have been detected in water, soil, food, and urine(3). In settings where glyphosate is applied, exposure to the chemical is highest. Still, outside of these settings, glyphosate is very common in the environment, contributing to long-term human exposure (3). Most non-occupational exposures occur through the consumption of contaminated foods, but may also be the result of contaminated soil, dust, and water (3). The food products that most commonly have detectable glyphosate residues include cereals and processed grain-based foods(1, 5). The herbicide enters surface water through a variety of routes, including run-off (5).

---

## Health Risks Associated with Glyphosate

### Cancer Health Risks

The International Agency for Research on Cancer (IARC) classifies glyphosate as a probable human carcinogen, due to sufficient evidence of cancer risk in experimental animals (5). In animal models, exposure to glyphosate has been associated with DNA damage and implicated in a process called oxidative stress (5). This occurs when our bodies have too many free radicals, which cause damage to human cells (5). While there is limited evidence in humans as to the carcinogenicity of glyphosate, the strongest evidence was for a relationship with non-Hodgkin lymphoma (NHL) (5). The relationship with other types of cancer was not as strong.

### Non-Cancer Health Risks

Glyphosate-based products can cause eye and skin irritation, and are potentially fatal when ingested (4). Because it is known that glyphosate interrupts cellular processes, more research is needed into adverse health outcomes associated with its use (5). Some studies have implicated glyphosate exposure in disease of the kidney, thyroid, and neurologic and reproductive systems; however, there is insufficient evidence to establish an association (6).

---

## How is Glyphosate regulated?

- The EPA establishes the maximum amounts of glyphosate on food products to range from 0.1 to 400 parts per million depending on the food product (7).
- The FDA monitors food and animal feed products for compliance with the EPA-established tolerances (8).
- In 2020, the FDA detected glyphosate in 54 of 2,800 human food samples (8).
- Glyphosate is not a restricted use pesticide, so no application certification or training is required for its use (4).

## What can you do?

- Limit personal use of glyphosate.
- Follow all safety precautions indicated on the product label (4).
- Wear personal protective equipment including long sleeves and pants, close-toed shoes, eye protection, and chemical resistant gloves when applying glyphosate-based herbicides (4).
- Allow 12 hours after application of glyphosate before re-entering the area or field (4).

*Published March 2025*

*Questions? Visit our website [EHSRC.org](http://EHSRC.org)*

# References



**IOWA**

College of Public Health

1. Environmental Protection Agency (2024). Glyphosate. <https://www.epa.gov/ingredients-used-pesticide-products/glyphosate>
2. Lacroix, R. & Kurrasch, D. M. (2023). Glyphosate toxicity: in vivo, in vitro, and epidemiological evidence. *Toxicological Sciences*, 192(2), pp. 131-140. <https://doi.org/10.1093/toxsci/kfad018>
3. Zhang, L., Rana, I., Shaffer, R. M., Taioli, E., & Sheppard, L. (2019). Exposure to glyphosate-based herbicides and risk for non-Hodgkin lymphoma: A meta-analysis and supporting evidence. *Mutation Research-Reviews in Mutation Research*, 781, pp. 186-206. doi: [10.1016/j.mrrev.2019.02.001](https://doi.org/10.1016/j.mrrev.2019.02.001)
4. Environmental Protection Agency (1993). R.E.D. Facts—Glyphosate. [https://www3.epa.gov/pesticides/chem\\_search/reg\\_actions/reregistration/fs\\_PC-417300\\_1-Sep-93.pdf](https://www3.epa.gov/pesticides/chem_search/reg_actions/reregistration/fs_PC-417300_1-Sep-93.pdf)
5. IARC Working Group on the Evaluation of Carcinogenic Risk to Humans (2017). IARC monographs on the evaluation of the carcinogenic risk of chemicals to man. Some organophosphate insecticides and herbicides. *IARC monographs on the evaluation of carcinogen risks to humans*, 112.
6. Agency for Toxic Substances and Disease Registry (2020). Toxicological Profile for Glyphosate. Centers for Disease Control and Prevention. <https://www.atsdr.cdc.gov/toxprofiles/tp214.pdf>
7. U.S. Food and Drug Administration. (2024 March 5). Questions and Answers on Glyphosate. <https://www.fda.gov/food/pesticides/questions-and-answers-glyphosate>
8. U.S. Food and Drug Administration (2022). Pesticide residue monitoring program fiscal year 2022 pesticide report. <https://www.fda.gov/media/181381/download?attachment>

# Nitrate & Health Fact sheet



**IOWA**

College of Public Health

---

## Introduction

Nitrate, a form of nitrogen, is a water-soluble chemical found at high concentrations in some Iowa waterways and private wells (1, 2, 3). Nitrate is colorless and has no smell, so it is unnoticeable in drinking water. Nitrate occurs naturally in the environment; however, applying nitrogen fertilizers to corn and other crop fields, and manure run-off from animal feeding operations can cause nitrate pollution in drinking water (2, 3). Nitrate can be reduced to nitrite and other compounds that have been shown to have negative health effects (1,5).

## Exposure Sources

People are primarily exposed to nitrate through contaminated drinking water and by eating certain nitrate-containing foods, like processed meats (1, 4). Drinking water contaminated with nitrate is a public health concern in Iowa due to the amount of nitrogen fertilizer used in agriculture (1, 2, 3), with shallow wells being most at risk of high nitrate concentrations (3). Nitrate is the most common form of nitrogen entering Iowa streams because of its stability and water-solubility, potentially contributing to downstream contamination of drinking water.

---

## Health Risks Associated with Nitrate

### Cancer Health Risks

The International Agency for Research on Cancer (IARC) classifies nitrate in food and water as "probably carcinogenic to humans" (1, 5). Studies conducted in Iowa have shown a positive correlation between exposure to nitrate in the drinking water and some cancers (1). Some studies conducted in Iowa have analyzed cancer risk at or above levels one-half the maximum contaminant level (MCL) over long periods of time. From these studies, consistent associations have been discovered for colorectal cancer. Cancers at other sites have been less studied, however, positive associations have been found at levels below the MCL for cancers of the thyroid, ovary, and kidney (6).

### Non-Cancer Health Risks

Methemoglobinemia, or Blue Baby Syndrome, is a condition in which not enough oxygen is transported in the blood to the body's tissues (3). Blue Baby Syndrome can occur when infants are exposed to nitrate levels higher than 10 mg/L. Elevated prenatal nitrate exposure in Iowa has also been connected to increased risk of birth defects, including spina bifida, limb deficiencies, and cleft palate (7,8).

---

## How is Nitrate regulated?

- The EPA established the maximum contaminant level for nitrate to be 10 mg/L in drinking water, to protect infants against methemoglobinemia, or Blue Baby Syndrome (7, 9).
- Public water suppliers must test for nitrate and present results to consumers and the state through consumer confidence reports (3).
- Private wells are not required to be tested for nitrate, and those with water sourced from a shallow well, less than 50 feet, face a disproportionate risk of high nitrate exposure (3,10).

## What can you do?

- Test well water for nitrate once a year if the well is privately-owned.
- Water purification methods that remove nitrate include reverse osmosis, ion exchange, and distillation. Using Brita filters, refrigerator filters, and boiling water are not effective at removing nitrate (7).
- Maintain a healthy diet rich in antioxidants and limited in processed, cured, and red meats (1). **Published September 2024**  
**Questions? Visit our website [EHSRC.org](https://www.ehsrc.org)**

# References



**IOWA**

College of Public Health

1. IARC Working Group on the Evaluation of Carcinogenic Risk to Humans (2010). IARC monographs on the evaluation of the carcinogenic risk of chemicals to man. Ingested nitrate and nitrite, and cyanobacterial peptide toxins. IARC monographs on the evaluation of carcinogen risks to humans, 94.
2. Iowa Water Quality Information System (n.d.) Water Quality Parameters Explained. From July 4, 2024. <https://iwqis.iowawis.org/parameters.html>
3. Iowa Department of Natural Resources (n.d.). Iowa Source Water Protection Guidebook. From July 4, 2024. <https://www.iowadnr.gov/Portals/idnr/uploads/water/wse/SWPPGuidebook.pdf>
4. Iowa Water Quality and Public Health Consortium (n.d.) Nitrate in Your Drinking Water: What You Need to Know. <https://ehsrc.public-health.uiowa.edu/wp-content/uploads/2021/01/Nitrates.pdf>
5. Grosse, Y., Baan, R., Straif, K., Secretan, B., El Ghissassi, F., Cogliano, V., & WHO International Agency for Research on Cancer Monograph Working Group (2006). Carcinogenicity of nitrate, nitrite, and cyanobacterial peptide toxins. *The Lancet. Oncology*, 7(8), 628–629. [https://doi.org/10.1016/s1470-2045\(06\)70789-6](https://doi.org/10.1016/s1470-2045(06)70789-6)
6. Ward, M. H., Jones, R. R., Brender, J. D., de Kok, T. M., Weyer, P. J., Nolan, B. T., Villanueva, C. M., & van Breda, S. G. (2018). Drinking Water Nitrate and Human Health: An Updated Review. *International journal of environmental research and public health*, 15(7), 1557. <https://doi.org/10.3390/ijerph15071557>
7. Iowa Environmental Council (2024). Nitrate in Drinking Water: A Public Health Concern for All Iowans. [https://www.iaenvironment.org/webres/File/IEC\\_Nitrate\\_in\\_Drinking\\_Water\\_2024FINAL.pdf](https://www.iaenvironment.org/webres/File/IEC_Nitrate_in_Drinking_Water_2024FINAL.pdf)
8. Brender, J. D., Weyer, P. J., Romitti, P. A., Mohanty, B. P., Shinde, M. U., Vuong, A. M., Sharkey, J. R., Dwivedi, D., Horel, S. A., Kantamneni, J., Huber, J. C., Jr, Zheng, Q., Werler, M. M., Kelley, K. E., Griesenbeck, J. S., Zhan, F. B., Langlois, P. H., Suarez, L., Canfield, M. A., & National Birth Defects Prevention Study (2013). Prenatal nitrate intake from drinking water and selected birth defects in offspring of participants in the national birth defects prevention study. *Environmental health perspectives*, 121(9), 1083–1089. <https://doi.org/10.1289/ehp.1206249>
9. United States Environmental Protection Agency (2024). National Primary Drinking Water Regulations. From July 4, 2024. <https://www.epa.gov/ground-water-and-drinking-water/national-primary-drinking-water-regulations>
10. Schechinger, Anne (2019). Iowa's Private Wells Contaminated by Nitrate and Bacteria. Environmental Working Group. From September 6, 2024. [https://www.ewg.org/interactive-maps/2019\\_iowa\\_wells/](https://www.ewg.org/interactive-maps/2019_iowa_wells/)

# Radon & Health Fact sheet



# IOWA

College of Public Health

## Introduction

Radon is a naturally occurring gas that originates from the breakdown of uranium and is present in air, water, and soil (1). It is odorless, colorless, tasteless, and imperceptible to the senses. Radon is radioactive, forming other radioactive products called decay products over time. Higher radon concentration in the ground in Iowa results from historical glacial activity that carried rock containing uranium to Iowa (2). The uranium in this rock naturally decays to form radon. Because the glaciers deposited this rock unevenly, uranium and, as a result, radon levels in the ground vary from site to site.

## Exposure Sources

Unstable radon decays into radioactive products and emits alpha particle radiation (1). These radioactive products can be inhaled and damage lungs the DNA in our lung cells. This damage can lead to genetic mutations, which may eventually cause lung cancer. Many people are exposed to radon by breathing in indoor air. The amount of radon in the air is measured in pCi/L (picocuries per liter of air). Radon gas can infiltrate homes from soil through small cracks in the foundation or openings around pipes, sump pumps, crawl spaces, drains, and other structures (2). Building materials, water supply, and natural gas are also sources of radon in the home.

Radon levels will increase in homes with poor ventilation and lower floors near the foundation, such as a basement or ground floor. Indoor radon levels are often higher than outdoors levels (1). Because of its unique glacial rock deposits, Iowa has the highest average indoor radon levels in the nation (2,3). Radon levels can vary from home to home because of the uneven distribution of radon in the ground (2). Radon testing is the only way to determine indoor radon concentrations (1,2,3).

## Health Risks Associated with Radon

### Cancer Health Risks

Radon causes cancer in humans (4,5) and, after smoking, is the second leading cause of lung cancer in the nation (1). For example, the Iowa Radon Lung Cancer Study concluded that “cumulative radon exposure in the residential environment is significantly associated with lung cancer risk” (5). The chances of getting lung cancer from radon increase with the indoor level of radon and the time spent in the home. People who smoke that are exposed to radon are at a 10 to 20 times higher risk of developing lung cancer than non-smokers (1). Moreover, with their different lung shapes and sizes and higher breathing rates, children are exposed to higher radon doses than adults. No consistent relationship has been found between elevated radon exposure and cancer at other sites.

### Non-Cancer Health Risks

Lung cancer is the only established adverse health effect associated with exposure to elevated radon levels (1). Some studies are being conducted in examination of a potential association between radon exposure and risk of stroke and clonal hematopoiesis of indeterminate potential (CHIP) (6).

## How is Radon regulated?

- There are no enforceable federal regulations for acceptable radon levels in indoor environments, such as individual residences or schools (1).
- The EPA recommends taking action to reduce radon levels if they are at or above 4 pCi/L to limit the risk of developing lung cancer (1).
- Radon levels below 4 pCi/L still carry risk and, in many cases, should be further reduced, especially for inhabited or heavily used areas (1).

## What can you do?

- Because the entire state of Iowa is at high risk of radon in homes and other buildings, the Iowa Department of Health and Human Services recommends testing every two years or after renovating your home (3). Radon testing is inexpensive and reliable.
- If the results from radon testing are greater than or equal to 2 pCi/L, installing a radon mitigation system can range from \$800 to \$2,500 (3).

*Updated September 2024 Questions? Visit our website [EHSRC.org](http://EHSRC.org)*

# References



1. Centers for Disease Control and Prevention (2023). Radon Toxicity: Standards and Regulations. Agency for Toxic Substances and Disease Registry. <https://www.atsdr.cdc.gov/csem/radon/standards.html>
2. Iowa Department of Health and Human Services (2020). Iowa Radon Home Buyers and Sellers Fact Sheet. <https://hhs.iowa.gov/media/7917/download?inline=#:~:text=Long%20ago%2C%20glacial%20activity%20left,harmful%20to%20humans%20when%20inhaled.>
3. Iowa Department of Health and Human Services (n.d.) Radon and You: What You Need to Know to Protect You and Your Family. [https://hhs.iowa.gov/media/7880/download?inline=.](https://hhs.iowa.gov/media/7880/download?inline=)
4. International Agency for Research on Cancer. (1988). Man-made mineral fibres and radon (IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, Vol. 43). <https://publications.iarc.fr/Book-And-Report-Series/Iarc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Man-Made-Mineral-Fibres-And-Radon-1988.>
5. Field, R. W., Steck, D. J., Smith, B. J., Brus, C. P., Fisher, E. L., Neuberger, J. S., Platz, C. E., Robinson, R. A., Woolson, R. F., Lynch, C. F. (2000). Residential Radon Gas Exposure and Lung Cancer: The Iowa Radon Lung Cancer Study. *American Journal of Epidemiology*, 151 (11), pp. 1091-1102. [https://cheec.uiowa.edu/sites/cheec.uiowa.edu/files/Residential%20Radon%20Gas%20Exposure%20and%20Lung%20Cancer\\_Iowa%20Study.pdf.](https://cheec.uiowa.edu/sites/cheec.uiowa.edu/files/Residential%20Radon%20Gas%20Exposure%20and%20Lung%20Cancer_Iowa%20Study.pdf)
6. UNC Gillings School of Global Public Health (2024) Is Radon Linked to a Health Condition Other Than Lung Cancer? From September 6, 2024. <https://sph.unc.edu/sph-news/is-radon-linked-to-a-health-condition-other-than-lung-cancer/>